

## A SHORT TABLE OF INDEFINITE INTEGRALS

### I. Basic Functions

- |   |  |
|---|--|
| <p>1. <math>\int x^n dx = \frac{1}{n+1}x^{n+1} + C, \quad n \neq -1</math></p> <p>2. <math>\int \frac{1}{x} dx = \ln x  + C</math></p> <p>3. <math>\int a^x dx = \frac{1}{\ln a}a^x + C</math></p> <p>4. <math>\int \ln x dx = x \ln x - x + C, \quad x &gt; 0</math></p> | <p>5. <math>\int \sin x dx = -\cos x + C</math></p> <p>6. <math>\int \cos x dx = \sin x + C</math></p> <p>7.(a) <math>\int \tan x dx = -\ln \cos x  + C</math></p> <p>7.(b) <math>\int \cot x dx = \ln \sin x  + C</math></p> <p>7.(c) <math>\int \sec^2 x dx = \tan x + C</math></p> <p>7.(d) <math>\int \csc^2 x dx = -\cot x + C</math></p> |
|---|--|

### II Products of $e^x$ , $\cos x$ , and $\sin x$

8.  $\int e^{ax} \sin(bx) dx = \frac{1}{a^2+b^2}e^{ax}[a \sin(bx) - b \cos(bx)] + C$
9.  $\int e^{ax} \cos(bx) dx = \frac{1}{a^2+b^2}e^{ax}[a \cos(bx) + b \sin(bx)] + C$
10.  $\int \sin(ax) \sin(bx) dx = \frac{1}{b^2-a^2}[a \cos(ax) \sin(bx) - b \sin(ax) \cos(bx)] + C, \quad a \neq b$
11.  $\int \cos(ax) \cos(bx) dx = \frac{1}{b^2-a^2}[b \cos(ax) \sin(bx) - a \sin(ax) \cos(bx)] + C, \quad a \neq b$
12.  $\int \sin(ax) \cos(bx) dx = \frac{1}{b^2-a^2}[b \sin(ax) \sin(bx) + a \cos(ax) \cos(bx)] + C, \quad a \neq b$

### III. Product of Polynomial $p(x)$ with $\ln x$ , $e^x$ , $\cos x$ , $\sin x$

13.  $\int x^n \ln x dx = \frac{1}{n+1}x^{n+1} \ln x - \frac{1}{(n+1)^2}x^{n+1} + C, \quad n \neq -1, \quad x > 0$
14.  $\int p(x)e^{ax} dx = \frac{1}{a}p(x)e^{ax} - \frac{1}{a} \int p'(x)e^{ax} dx$   
 $= \frac{1}{a}p(x)e^{ax} - \frac{1}{a^2}p'(x)e^{ax} + \frac{1}{a^3}p''(x)e^{ax} - \dots$   
 (+ - + - ...) (signs alternate)
15.  $\int p(x) \sin ax dx = -\frac{1}{a}p(x) \cos ax + \frac{1}{a} \int p'(x) \cos ax dx$   
 $= -\frac{1}{a}p(x) \cos ax + \frac{1}{a^2}p'(x) \sin ax + \frac{1}{a^3}p''(x) \cos ax - \dots$   
 (- + + - - + + ...) (signs alternate in pairs after first term)
16.  $\int p(x) \cos ax dx = \frac{1}{a}p(x) \sin ax - \frac{1}{a} \int p'(x) \sin ax dx$   
 $= \frac{1}{a}p(x) \sin ax + \frac{1}{a^2}p'(x) \cos ax - \frac{1}{a^3}p''(x) \sin ax - \dots$   
 (+ + - - + + - - ...) (signs alternate in pairs)

#### IV. Integer Powers of $\sin x$ and $\cos x$

$$17. \int \sin^n x dx = -\frac{1}{n}(\sin^{n-1} x) \cos x + \frac{n-1}{n} \int \sin^{n-2} x dx, \quad n \text{ positive}$$

$$18. \int \cos^n x dx = \frac{1}{n}(\cos^{n-1} x) \sin x + \frac{n-1}{n} \int \cos^{n-2} x dx, \quad n \text{ positive}$$

$$19. \int \frac{1}{\sin^m x} dx = \frac{-1}{(m-1)} \frac{\cos x}{\sin^{m-1} x} + \frac{m-2}{m-1} \int \frac{1}{\sin^{m-2} x} dx, \quad m \neq 1, \quad m \text{ positive}$$

$$20. \int \frac{1}{\sin x} dx = \int \csc x dx = \ln[\csc x - \cot x] + C$$

$$21. \int \frac{1}{\cos^m x} dx = \frac{1}{(m-1)} \frac{\sin x}{\cos^{m-1} x} + \frac{m-2}{m-1} \int \frac{1}{\cos^{m-2} x} dx, \quad m \neq 1, \quad m \text{ positive}$$

$$22. \int \frac{1}{\cos x} dx = \int \sec x dx = \ln[\sec x + \tan x] + C$$

$$23. \int \sin^m x \cos^n x dx : \text{ If } m \text{ is odd, let } w = \cos x. \text{ If } n \text{ is odd, let } w = \sin x.$$

If both  $m$  and  $n$  are even and non-negative, convert all to  $\sin x$  or all to  $\cos x$  (using  $\sin^2 x + \cos^2 x = 1$ ), and use IV-17 or IV-18. If  $m$  and  $n$  are even and one of them is negative, convert to whichever function is in the denominator and use IV-19 or IV-21. The case in which both  $m$  and  $n$  are even and negative is omitted.

#### V. Quadratic in the Denominator

$$24. \int \frac{1}{x^2+a^2} dx = \frac{1}{a} \arctan \frac{x}{a} + C, \quad a \neq 0$$

$$25. \int \frac{bx+c}{x^2+a^2} dx = \frac{b}{2} \ln|x^2+a^2| + \frac{c}{a} \arctan \frac{x}{a} + C, \quad a \neq 0$$

$$26. \int \frac{1}{(x-a)(x-b)} dx = \frac{1}{a-b} (\ln|x-a| - \ln|x-b|) + C, \quad a \neq b$$

$$27. \int \frac{cx+d}{(x-a)(x-b)} dx = \frac{1}{a-b} [(ac+d) \ln|x-a| - (bc+d) \ln|x-b|] + C, \quad a \neq b$$

#### VI: Integrands involving $\sqrt{a^2+x^2}$ , $\sqrt{a^2-x^2}$ , $\sqrt{x^2-a^2}$ , $a > 0$

$$28. \int \frac{1}{\sqrt{a^2-x^2}} dx = \arcsin \frac{x}{a} + C$$

$$29. \int \frac{1}{\sqrt{x^2 \pm a^2}} dx = \ln|x + \sqrt{x^2 \pm a^2}| + C$$

$$30. \int \sqrt{a^2 \pm x^2} dx = \frac{1}{2} \left( x\sqrt{a^2 \pm x^2} + a^2 \int \frac{1}{\sqrt{a^2 \pm x^2}} dx \right) + C$$

$$31. \int \sqrt{x^2 - a^2} dx = \frac{1}{2} \left( x\sqrt{x^2 - a^2} - a^2 \int \frac{1}{\sqrt{x^2 - a^2}} dx \right) + C$$