

MATHEMATICS 271

MIDTERM COVER PAGE

March 13, 2008

NAME_____Lecture Section/Professor_____

NAME _____ ID _____ Section _____

MATHEMATICS 271 MIDTERM March 13, 2008

SHOW ALL WORK. Marks for each problem are to the left of the problem number.
NO CALCULATORS PLEASE.

- [6] 1. Use the Euclidean algorithm to find $\gcd(72, 17)$. Then use your work to write $\gcd(72, 17)$ in the form $72a + 17b$ where a and b are integers.

- [6] 2. Let \mathcal{S} be the statement:

for all sets A and B , if $A \cup B = \{1, 2\}$ then $\{1, 2\} \in \mathcal{P}(A) \cup \mathcal{P}(B)$.

(Here $\mathcal{P}(X)$ denotes the power set of the set X .)

- (a) Write (as simply as possible) the *negation* of statement \mathcal{S} .

- (b) *Disprove* statement \mathcal{S} .

ID_____

[11] 3. Let \mathcal{S} be the statement:

for all integers n , if $6 \mid n$ then $9 \mid (n^2 + 3n)$.

(a) Is \mathcal{S} true? Give a proof or disproof.

(b) Write out (as simply as possible) the *contrapositive* of statement \mathcal{S} . Is it true or false?
Explain.

(c) Write out (as simply as possible) the *converse* of statement \mathcal{S} . Is it true or false?
Explain.

ID_____

[6] 4. Of the following two statements, one is true and one is false. Prove the true statement. Disprove the false statement by writing out its negation and proving that. (\mathbb{Z} denotes the set of all integers.)

(a) $\forall A \subseteq \mathbb{Z} \exists B \subseteq \mathbb{Z}$ so that $1 \in B - A$.

(b) $\forall A \subseteq \mathbb{Z} \exists B \subseteq \mathbb{Z}$ so that $1 \notin B - A$.

[5] 5. You are given that A and B are arbitrary subsets of the set \mathbb{Z} of all integers such that $A \cap B = \{1\}$.

(a) Find an element of $A \times B$. Explain.

(b) Find an element of the complement $(A \times B)^c$. (Here assume the universal set is $\mathbb{Z} \times \mathbb{Z}$.) Explain.

ID_____

[6] 6. Prove **using mathematical induction** (or well ordering) that $2^n + 2n \leq 3^n$ for all integers $n \geq 2$.