

1. Determine whether or not the following sets with the indicated operations are vector spaces or not. Justify your answer.
- The set, V , of non-negative real numbers; with ordinary addition and scalar multiplication.
 - The set, V , of all 2×2 matrices of the form $\begin{pmatrix} a & b \\ 0 & c \end{pmatrix}$; with the usual operations of addition and scalar multiplication as for the vector space M_{22} .
 - The set, V , of all 2×2 matrices which have determinant equal to zero; with the usual operations of addition and scalar multiplication as for the vector space M_{22} .
 - The set V of all 2×2 matrices with the addition of M_{22} but having scalar multiplication, $*$, defined by: for $a \in \mathbb{R}$, and $X \in V$, $a * X = aX^T$
 - Let V be a set which consists of the ordered triples (x, y, z) in \mathbb{R}^3 with addition defined as usual and with scalar multiplication defined as follows:
If $a \in \mathbb{R}$, and $(x, y, z) \in \mathbb{R}^3$, then $a(x, y, z) = (ax, 0, az)$
 - Let V be the set of all ordered pairs, $\begin{pmatrix} x \\ y \end{pmatrix}$, in \mathbb{R}^2 with scalar multiplication defined as usual and addition defined as follows:

$$\text{If } \begin{pmatrix} x_1 \\ y_1 \end{pmatrix}, \begin{pmatrix} x_2 \\ y_2 \end{pmatrix} \in \mathbb{R}^2, \text{ then } \begin{pmatrix} x_1 \\ y_1 \end{pmatrix} + \begin{pmatrix} x_2 \\ y_2 \end{pmatrix} = \begin{pmatrix} x_1 + x_2 - 1 \\ y_1 + y_2 \end{pmatrix}$$

2. Let V be the set of positive real numbers with vector addition being ordinary multiplication, and scalar multiplication being defined by $a \mathbf{v} = \mathbf{v}^a$. Show that V is a vector space.

University of Calgary
Faculty of Science
Department of Mathematics and Statistics

Math 311

Sheet 2

Spring 2008

3. In each case show that the condition $au + bv + cw = \mathbf{0}$ in V implies that $a = b = c = 0$.
- a. $V = \mathbb{R}^4$; $u = (2,1,0,2)$, $v = (1,1,-1,0)$, $w = (0,1,2,1)$
- b. $V = P_3$; $u = x^3 + x$, $v = x^2 + 1$, $w = x^3 - x^2 + x + 1$
4. Let V be a vector space which contains the vectors v, v_1, v_2, \dots, v_n . Let a_1, a_2, \dots, a_n be real numbers. Use induction on n to prove that
- a. $a(v_1 + v_2 + \dots + v_n) = av_1 + av_2 + \dots + av_n$
- b. $(a_1 + a_2 + \dots + a_n)v = a_1v + a_2v + \dots + a_nv$
5. Let V be a vector space. If u and v are vectors in V , and if a and b are scalars, show that
- a. If $av = bv$ and $v \neq \mathbf{0}$, then $a = b$
- b. If $au = av$ and $a \neq 0$, then $u = v$