

**The University of Calgary**  
**Department of Mathematics and Statistics**  
**MATH 349      Handout # 5**

1. For  $f(x, y) = \frac{xy}{\sqrt{1+x^2}}$  and  $\mathbf{g} : \mathbb{R}^2 \rightarrow \mathbb{R}^2, \mathbf{g}(s, t) = (\cos(\pi st), \sin \frac{\pi s}{t})$ 
  - (a) find  $\nabla f$ ;
  - (b) using Chain Rule find  $\nabla h(0, -1)$  where  $h = f \circ \mathbf{g}$   
 ( or  $h(s, t) = f(x, y)$  where  $x = \cos(\pi st)$  and  $y = \sin \frac{\pi s}{t}$ )
  
2. Find an equation of the tangent plane to  $z = f(x, y) = \ln(x + y^2)$  at the point  $x_0 = 0, y_0 = -1$ .
  
3. For the function  $f(x, y) = e^{\sqrt{\frac{y}{x}}}$  find the domain and  $f_{xx}$  and  $f_{xy}$ .
  
4. For  $f(x, y, z) = \sqrt{2} \sin(\pi xy + x \ln z)$  and  $\mathbf{g} : \mathbb{R} \rightarrow \mathbb{R}^3, \mathbf{g}(t) = (\frac{1}{t}, -\frac{1}{t}, \frac{t}{2})$ 
  - (a) find  $\nabla f$ ;
  - (b) find  $D\mathbf{g}$  or  $\mathbf{g}'$
  - (c) using Chain Rule find  $h'(2)$  where  $h = f \circ \mathbf{g}$   
 ( or  $h(t) = f(x, y, z)$  where  $x = \frac{1}{t}, y = -\frac{1}{t}$  and  $z = \frac{t}{2}$ )
  
5. In what directions at the point  $P(2, 1)$  does the function  $f(x, y) = \ln(\frac{x}{y} + \frac{y}{x})$  have the rate of change equal to  $\frac{3}{10}$ ? What is the maximum rate of change at that point?
  
6. Find the rate and the direction of the most rapid decrease of  $f(x, y, z) = x^2 z e^y + x z^2$  at the point  $P(1, \ln 2, 2)$ .
  
7. Given  $f(x, y) = \begin{cases} \frac{xy}{2x^2 + y^2} & \text{for } (x, y) \neq (0, 0) \\ 0 & \text{.....at } (0, 0) \end{cases}$ 
  - (a) Is  $f$  continuous at  $(0, 0)$ ?
  - (b) Find  $\nabla f$  at  $(0, 0)$ , if it exists.
  - (c) Find the directional derivative at  $(0, 0)$  in the direction of  $y = x$  if it exists.
  - (d) Find the directional derivative at  $(-1, -1)$  where  $\mathbf{u} = \frac{1}{\sqrt{2}}(1, 1)$ .