

The University of Calgary
Faculty of Science
Department of Mathematics and Statistics
MATH 353
Midterm Supplement

1. Describe in the xyz - space the following sets

(a) where $\rho = x$ (ρ is from spherical coord.);

(b) where $\rho = -2y$.

for 1a)

$$\sqrt{x^2 + y^2 + z^2} = x \text{ so } x \geq 0 \text{ and } x^2 + y^2 + z^2 = x^2, y^2 + z^2 = 0$$

thus $y = z = 0$ and any point $(x, 0, 0)$ for $x \geq 0$ - half of the x-axis

for b)

$$\sqrt{x^2 + y^2 + z^2} = -2y \text{ so } y \leq 0 \text{ and } x^2 + y^2 + z^2 = 4y^2$$

$$x^2 + z^2 = 3y^2, \text{ one part cone around the } y\text{-axis} \quad y = -\frac{\sqrt{x^2 + z^2}}{\sqrt{3}}.$$

2. Set up the integral $\iiint_B z \, dx \, dy \, dz$ where B is the region in the first octant below the plane

$$z = 2$$

and above the plane $3x + 2y - 6z = 0$ as iterated integrals

(a) with $\int dz$ inside ;

(b) with $\int dz$ outside

then evaluate only once.

for 2a)

$$\text{first } B = \{(x, y, z); x \geq 0, y \geq 0, z \geq 0, \frac{3x + 2y}{6} \leq z \leq 2\}$$

"bottom" must be below "top" $\frac{3x + 2y}{6} \leq 2$ and we get

$$D = \{(x, y); x \geq 0, y \geq 0, 3x + 2y \leq 12\} \quad \text{a triangle}$$

$$\text{so } \iiint_B z \, dx \, dy \, dz = \iint_D \left(\int_{\frac{3x+2y}{6}}^2 z \, dz \right) dx \, dy = \int_0^4 \left[\int_0^{\frac{12-3x}{2}} \left(\int_{\frac{3x+2y}{6}}^2 z \, dz \right) dy \right] dx = ..$$

for b)

$0 \leq z \leq 2$ and for a fixed z

$$D_z = \{(x, y); x \geq 0, y \geq 0, 3x + 2y \leq 6z\}$$

triangles with the base $[0, 2z]$ and the height $[0, 3z]$ and the area $3z^2$

$$\iiint_B z \, dx \, dy \, dz = \int_0^2 z \left(\iint_{D_z} dx \, dy \right) dz = \int_0^2 z (\text{area of } D_z) dz = 3 \int_0^2 z^3 dz = 12$$

OR

$$\iiint_B z \, dx dy dz = \int_0^2 z \left(\int_0^{2z} \left(\int_0^{\frac{6z-3x}{2}} dy \right) dx \right) dz = \dots$$

3. Set up the integral $\iiint_B \frac{dx dy dz}{\sqrt{x^2 + y^2 + z^2}}$

where $B = \{(x, y, z); x^2 + y^2 + z^2 \leq 4; x^2 + y^2 \geq 3, x \geq 0, y \geq 0\}$
as iterated integrals

- (a) in spherical coordinates;
- (b) in cylindrical coordinates,
then evaluate only once.

for 3 a)

the solid B is outside the cylinder and inside the sphere

$$x \geq 0, y \geq 0 \text{ implies } \theta \in [0, \frac{\pi}{2}], x^2 + y^2 + z^2 \leq 4 \rightarrow 0 \leq \rho \leq 2$$

$$\text{and } x^2 + y^2 \geq 3 \text{ implies } \rho \sin \phi \geq \sqrt{3} \quad \frac{\sqrt{3}}{\sin \phi} \leq \rho \leq 2$$

$$\text{so necessarily } \frac{\sqrt{3}}{2} \leq \sin \phi \quad \phi \in \left[\frac{\pi}{3}, \frac{2\pi}{3} \right]$$

$$B^* = \left\{ \theta \in [0, \frac{\pi}{2}], \phi \in \left[\frac{\pi}{3}, \frac{2\pi}{3} \right], \frac{\sqrt{3}}{\sin \phi} \leq \rho \leq 2 \right\}$$

as iterated integrals

$$\iiint_B \frac{dx dy dz}{\sqrt{x^2 + y^2 + z^2}} = \iiint_{B^*} \frac{\rho^2 \sin \phi d\rho d\phi d\theta}{\rho} = \frac{\pi}{2} \int_{\frac{\pi}{3}}^{\frac{2\pi}{3}} \sin \phi \left(\int_{\frac{\sqrt{3}}{\sin \phi}}^2 \rho \, d\rho \right) d\phi =$$

$$= \frac{\pi}{4} \int_{\frac{\pi}{3}}^{\frac{2\pi}{3}} \sin \phi \left(4 - \frac{3}{\sin^2 \phi} \right) d\phi = \pi \left[-\cos \phi \right]_{\frac{\pi}{3}}^{\frac{2\pi}{3}} - \frac{3\pi}{4} \left[\ln |\csc \phi - \cot \phi| \right]_{\frac{\pi}{3}}^{\frac{2\pi}{3}} =$$

$$\left(\cos \frac{\pi}{3} = \frac{1}{2}, \cos \frac{2\pi}{3} = -\frac{1}{2}, \sin(\text{both}) = \frac{\sqrt{3}}{2}, \cot = \pm \frac{1}{\sqrt{3}} \right)$$

$$= \pi - \frac{3\pi}{2} \ln \sqrt{3} = \pi \left(1 - \frac{3}{4} \ln 3 \right)$$

for b)

the solid B is outside the cylinder and inside the sphere

$$x \geq 0, y \geq 0 \text{ implies } \theta \in [0, \frac{\pi}{2}], x^2 + y^2 + z^2 \leq 4 \rightarrow r^2 + z^2 \leq 4$$

$$\text{and } x^2 + y^2 \geq 3 \text{ implies } r \geq \sqrt{3}$$

$$B^* = \left\{ \theta \in [0, \frac{\pi}{2}], r \geq \sqrt{3}, r^2 + z^2 \leq 4 \right\}$$

$$\iiint_B \frac{dx dy dz}{\sqrt{x^2 + y^2 + z^2}} = \iiint_{B^*} \frac{r dr dz d\theta}{\sqrt{r^2 + z^2}} = \frac{\pi}{2} \int_{\sqrt{3}}^2 r \left(\int_{-\sqrt{4-r^2}}^{\sqrt{4-r^2}} \frac{dz}{\sqrt{r^2 + z^2}} \right) dr =$$

$$\begin{aligned} \text{OR} &= \frac{\pi}{2} \int_{-1}^1 \left(\int_{\sqrt{3}}^{\sqrt{4-z^2}} \frac{r \, dr}{\sqrt{r^2+z^2}} \right) dz = \pi \int_0^1 \left(\sqrt{r^2+z^2} \right)_{\sqrt{3}}^{\sqrt{4-z^2}} dz = \pi \int_0^1 (2 - \sqrt{3+z^2}) dz = \\ &= 2\pi - \frac{\pi}{2} \left[z\sqrt{3+z^2} + 3 \ln \left| z + \sqrt{3+z^2} \right| \right]_0^1 = 2\pi - \frac{\pi}{4} [2 + 3 \ln 3 - 3 \ln \sqrt{3}] = \\ &= \pi \left[1 - \frac{3}{4} \ln 3 \right]. \end{aligned}$$

4. Evaluate $\iiint_B \frac{dx dy dz}{z-6}$ where B is the solid bounded by planes $x=0, y=0, z=0,$

and $3x+3y+z=6$. HINT: Iterate in such a way that $\int dz$ is outside!

for 4)

the solid is a tetrahedron

in the first octant under the plane $3x+3y+z=6$

so $B = \{x \geq 0, y \geq 0, z \geq 0, 3x+3y+z \leq 6\}$

for $z=0$ we have a triangle $(0,0), (2,0), (0,2)$

for $z=6$ $x=y=0$

for $z \in (0,6)$ we have a smaller triangle

under the line $3x+3y=6-z$

with vertices at $(0,0), (0, \frac{6-z}{3}), (\frac{6-z}{3}, 0)$

$D_z = \{x \geq 0, y \geq 0, x+y \leq \frac{6-z}{3}\}$...triangle with area $A = \frac{1}{2} \left(\frac{6-z}{3}\right)^2$

$$\begin{aligned} \iiint_B \frac{dx dy dz}{z-6} &= \int_0^6 \frac{1}{z-6} \left(\iint_{D_z} dx dy \right) dz = \int_0^6 \frac{1}{z-6} \cdot \text{area} dz = \frac{1}{18} \int_0^6 (z-6) dz = \\ &= \frac{1}{36} [(z-6)^2]_0^6 = \frac{-36}{36} = -1. \end{aligned}$$

5. Set the integral $\iiint_B z \, dx dy dz$ where $B = \{x^2+y^2+z^2 \leq 2, z \geq 0, y \geq 0, x^2+y^2 \geq z\}$

as iterated integrals in both

(a) cylindrical and

(b) spherical coordinates then evaluate only one of the above .

for 5a)

the set is inside the sphere with radius $\sqrt{2}$

below the paraboloid $z = x^2 + y^2$, and above the xy-plane $z = 0$

$B^* = \{r^2 + z^2 \leq 2, z \geq 0, r \geq 0, \theta \in [0, \pi], r^2 \geq z\}$

$$\text{so } I = \iiint_B z \, dx dy dz = \iiint_{B^*} z \, r dr d\theta dz = \pi \iint_D r z \, dr dz$$

where $D = \{r^2 + z^2 \leq 2, z \geq 0, r \geq 0, r^2 \geq z\}$

intersection of the circle and parabola is at $z = r = 1$

it is easier to slice it horizontally in zr - plane

so $z \in [0, 1]$ and $r \geq \sqrt{z}$, $r \leq \sqrt{2-z^2}$

$$\text{the integral } I = \pi \int_0^1 z \left(\int_{\sqrt{z}}^{\sqrt{2-z^2}} r dr \right) dz = \pi \int_0^1 z \left(\left[\frac{r^2}{2} \right]_{\sqrt{z}}^{\sqrt{2-z^2}} \right) dz =$$

$$= \frac{1}{2} \pi \int_0^1 z (2 - z^2 - z) dz = \frac{\pi}{2} \left[z^2 - \frac{z^4}{4} - \frac{z^3}{3} \right]_0^1 = \frac{\pi}{2} \cdot \left(1 - \frac{7}{12} \right) = \frac{5}{24} \pi.$$

For b)

since $x^2 + y^2 = \rho^2 \sin^2 \phi$

$$B^* = \left\{ \rho \leq \sqrt{2}, \theta \in [0, \pi], \phi \in [0, \frac{\pi}{2}], \rho \geq \frac{\cos \phi}{\sin^2 \phi} \right\}$$

but necessarily $\sqrt{2} \geq \frac{\cos \phi}{\sin^2 \phi} \rightarrow \phi \in [\frac{\pi}{4}, \frac{\pi}{2}]$

thus

$$I = \iiint_B z dx dy dz = \iiint_{B^*} \rho^3 \cos \phi \sin \phi d\rho d\phi d\theta =$$

$$= \int_0^\pi d\theta \cdot \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \left(\cos \phi \sin \phi \int_{\frac{\cos \phi}{\sin^2 \phi}}^{\sqrt{2}} \rho^3 d\rho \right) d\phi \dots$$

6. Evaluate $\iiint_B \sqrt{x^2 + 2y^2} dx dy dz$ where B is the solid bounded

by surfaces $z = x^2 + y^2$ and $z = 4 - x^2 - 3y^2$.

For 6)

$B = \{x^2 + y^2 \leq z \leq 4 - x^2 - 3y^2\}$ between two paraboloids

necessarily $x^2 + y^2 \leq 4 - x^2 - 3y^2$

so $2x^2 + 4y^2 \leq 4$ that is $(x, y) \in D_0 = \{x^2 + 2y^2 \leq 2\}$

and

$$\iiint_B \sqrt{x^2 + 2y^2} dx dy dz = \iint_{D_0} \sqrt{x^2 + 2y^2} \left(\int_{x^2+y^2}^{4-x^2-3y^2} dz \right) dx dy =$$

$$= \iint_{D_0} \sqrt{x^2 + 2y^2} (4 - 2(x^2 + 2y^2)) dx dy$$

use modified polar coord. to evaluate $x = \sqrt{2}r \cos \theta, y = r \sin \theta$

then $x^2 + 2y^2 = 2r^2$ $dx dy = \sqrt{2}r dr d\theta$

so D_0 transforms into $D^* = \{0 \leq r \leq 1, 0 \leq \theta \leq 2\pi\}$

and the integral into

$$= \int_0^{2\pi} d\theta \cdot 4\sqrt{2} \int_0^1 (r - r^3) \sqrt{2}r dr = 8 \cdot 2\pi \left[\frac{r^3}{3} - \frac{r^5}{5} \right]_0^1 = 16\pi \cdot \frac{2}{15} = \frac{32}{15} \pi.$$

7. For the solid B in the first octant bounded by the coordinate planes, the plane $y + z = 2$ and the surface $x = 4 - y^2$.

set up two different ways of integration of $\iiint_B f dx dy dz$

(a) (double first, then single) $\iint_{D_0} \left(\int f dz \right) dx dy$; sketch D_0 ;

(b) (single first, then double) $\int_a^b \left(\iint_{D_z} f dx dy \right) dz$; sketch D_z .

For 7)

$$B = \{x \geq 0, y \geq 0, z \geq 0, z \leq 2 - y, x \leq 4 - y^2 \text{ (or } y \leq \sqrt{4 - x} \text{)}\}$$

so

$$D_0 = \{x \geq 0, y \geq 0, x \leq 4 - y^2 \text{ (or } y \leq \sqrt{4 - x} \text{)}\}$$

and for a)

$$\iiint_B f dx dy dz = \iint_{D_0} \left(\int_0^{2-y} f dz \right) dx dy$$

for b)

$$0 \leq z \leq 2 \text{ and for a fixed } z \quad D_z = \{0 \leq y \leq 2 - z, 0 \leq x \leq 4 - y^2\}$$

$$\iiint_B f dx dy dz = \int_0^2 \left(\iint_{D_z} f dx dy \right) dz$$

