## MATH 353 Handout #2

- 1. Find absolute extrema of  $f(x,y) = \frac{1}{8}x^3 + y^3$  on the circle  $x^2 + y^2 \le 65$
- 2. Find the absolute extrema of  $f(x,y) = x^2 + y^2$  on the surface  $S = \left\{ \frac{1}{8}x^3 + y^3 = 65, x \ge 0, y \ge 0. \right\}$ .
- 3. Find absolute maxim and minima of  $f(x,y) = 2y^2 x + x^2$  inside and on the triangle T with vertices O(0,0), A(1,1), B(1,-1).
- 4. Find the point on the plane x-2y-z=3 closest to the point  $P\left(1,-1,2\right)$ . Justify!
- 5. Find absolute maximum of f(x, y, z) = xyz for  $x, y, z \ge 0$  on the surface 2xy + 2xz + 3yz = 144. (You may assume that there is an absolute maximum).
- 6. (a) Evaluate  $\int_{1}^{3} \left( \int_{-x}^{x^2} xe^{2y} dy \right) dx$ .
  - (b) Switch the order of integration in the integral above and sketch the region D.
- 7. Evaluate  $\iint_D \sqrt{2-x^2} dA$  where D is smaller region between  $y=x^2$  and  $x^2+y^2=2$ . and sketch the region
- 8. Switch the order of integration in the integral  $\int_{0}^{\frac{\pi}{4}} \left( \int_{0}^{\tan x} f(x,y) dy \right) dx$ .
- 9. For  $\iint_D \frac{1}{x^2 + y} dA$  where D is the region between the x-axis and  $y = 4 x^2$  sketch the region D and set up BOTH iterated integrals and evaluate one of them. (Hint:  $\lim_{x \to 0^+} x \ln x = 0$ ).
- 10. Calculate the volume of the solid below the surface  $z = e^{(y-1)^2}$  and above the triangle T with vertices

$$A(-1,0)$$
,  $B(0,1)$ ,  $C(2,0)$  with vertical sides.