

PMAT 421      WINTER 99      FINAL  
 3 hours  
 SOLUTIONS

**For 1)**

$$\left(-\frac{\pi}{2}\right)^{\frac{1}{i}} = e^{\frac{1}{i} \log\left(\frac{-\pi}{2}\right)} = e^{\frac{1}{i} [\ln \frac{\pi}{2} + i(\pi + 2k\pi)]} = e^{-i \ln \frac{\pi}{2}} \cdot e^{\pi(2k+1)} = e^{\pi(2k+1)} \cos\left(\ln \frac{\pi}{2}\right) - ie^{\pi(2k+1)} \sin\left(\ln \frac{\pi}{2}\right) \text{ for any integer } k.$$

**For 2)**

we can use the definition of  $\sin : \frac{e^{iz} - e^{-iz}}{2i} = 2$  and  $w = e^{iz}$

we get  $w - \frac{1}{w} = 4i$  and then  $w^2 - 4iw - 1 = 0$

$$w_{1/2} = \frac{4i \pm \sqrt{-16+4}}{2} = i(2 \pm \sqrt{3}) \text{ or } w_1 = i(2 + \sqrt{3}), w_2 = i(2 - \sqrt{3}) = \frac{i}{2 + \sqrt{3}}$$

now

$$z = \frac{1}{i} \log w = -i \left[ \ln(2 \pm \sqrt{3}) + i\left(\frac{\pi}{2} + 2k\pi\right) \right] = \frac{\pi}{2} + 2k\pi \mp i \ln(2 + \sqrt{3})$$

for any integer  $k$ .

**For 3)**

the property of  $\log : \log w^2 = 2 \log w + i2n\pi$  for certain  $n$  ONLY

it means that  $2 \log(-z) = 2 \log z + i2n\pi$

in detail if  $|\text{Arg } z| > \frac{\pi}{2}$  i.e. the point  $z$  is in the left half

$2 \cdot |\text{Arg } z| > \pi$  so it is not principal value anymore thus in this case

$$\text{Arg } z^2 \neq 2 \cdot \text{Arg } z.$$

**For 4)**

the center is 1 and singular point is 4 so we have two possible domains

$|z - 1| < 3$  or  $|z - 1| > 3$  we need the latter so negative powers of  $(z - 1)$  :

first

$$\frac{1}{z-4} = \frac{1}{(z-1)-3} = \frac{1}{z-1} \cdot \frac{1}{1-\frac{3}{z-1}} = \left( \text{for } \left| \frac{3}{z-1} \right| < 1 \right)$$

$$= \frac{1}{z-1} \sum_{n=0}^{\infty} \frac{3^n}{(z-1)^n} = \sum_{n=0}^{\infty} \frac{3^n}{(z-1)^{n+1}} = \sum_{k=1}^{\infty} 3^{k-1} (z-1)^{-k}$$

differentiate

$$\frac{-1}{(z-4)^2} = \sum_{k=1}^{\infty} 3^{k-1} (-k) (z-1)^{-k-1} \quad \text{and for } |z-1| > 3$$

$$\frac{1}{(z-4)^2} = \sum_{k=1}^{\infty} 3^{k-1} (k) (z-1)^{-k-1} = \sum_{k=1}^{\infty} \frac{k3^{k-1}}{(z-1)^{k+1}} = \sum_{n=2}^{\infty} \frac{(n-1)3^{n-2}}{(z-1)^n}.$$

**For 5)**

the function  $\cos$  is  $2\pi$ -periodic so if we can find one  $z$  we have infinitely many.

By contradiction:

if there is no such  $z$  it means that  $|\cos z| \leq 100$  for all  $|z| > R$

but  $\cos$  is continuous for  $|z| \leq R$  so  $|\cos z| \leq M$  for some  $M > 0$

together

$\cos$  is entire and bounded so by Liouville's Th.  $\cos z = \text{const.}$  which is NOT true.

Liouville's Th.

If  $f$  is entire and bounded

i.e.  $f'(z)$  exists for all  $z$  and for some  $M > 0$   $|f'(z)| \leq M$  for all  $z$ ;

then  $f$  is constant .

**For 6)**

since  $\text{Im } z$  is nowhere analytic we have to use the definition of the integral

first parametrize the curve  $c : |z - i| = 1$

$$z(t) = i + e^{it}, t \in \left[-\frac{\pi}{2}, 0\right], \text{Im } z(t) = 1 + \sin t, dz = ie^{it} dt$$

$$\begin{aligned} \int_c \text{Im } z dz &= \int_{-\frac{\pi}{2}}^0 (1 + \sin t) i (\cos t + i \sin t) dt = \\ &= i \int_{-\frac{\pi}{2}}^0 [\cos t + \cos t \sin t] dt - \int_{-\frac{\pi}{2}}^0 (\sin t + \sin^2 t) dt = \\ &= \left[ i \sin t + i \frac{\sin^2 t}{2} + \cos t \right]_{-\frac{\pi}{2}}^0 - \int_{-\frac{\pi}{2}}^0 \frac{1 - \cos 2t}{2} dt = 1 + i - \frac{i}{2} - \frac{1}{2} \cdot \frac{\pi}{2} + \left[ \frac{\sin 2t}{4} \right]_{-\frac{\pi}{2}}^0 = \\ &= \left(1 - \frac{\pi}{4}\right) + \frac{1}{2}i. \end{aligned}$$

**For 7a)**

$z \sin z = 0$  for  $z = k\pi$  for any integer  $k$

for  $k = 0$   $z_0 = 0$  is a pole of order  $m = 2$  since  $\lim_{z \rightarrow 0} z^2 f(z) = \lim_{z \rightarrow 0} \frac{z}{\sin z} = 1 \neq 0$

for  $k \neq 0$   $z_k = k\pi$  are simple poles

$$\text{since } \lim_{z \rightarrow k\pi} (z - k\pi) f(z) = \frac{1}{k\pi} \cdot \lim_{z \rightarrow k\pi} \frac{z - k\pi}{\sin z} = \frac{1}{k\pi \cos k\pi} = \frac{(-1)^k}{k\pi} \neq 0$$

**for b)**

$$\begin{aligned} \text{Res}[f, 0] &= \lim_{z \rightarrow 0} [z^2 f(z)]' = \lim_{z \rightarrow 0} \left[ \frac{z}{\sin z} \right]' = \lim_{z \rightarrow 0} \frac{\sin z - z \cos z}{\sin^2 z} (L'H) = \\ &= \lim_{z \rightarrow 0} \frac{\cos z - \cos z + z \sin z}{2 \sin z \cos z} = 0. \end{aligned}$$

**For 8)**

$$\int_0^\infty \frac{x^2}{x^4 + 4} dx = \frac{1}{2} \int_{-\infty}^\infty \frac{x^2}{x^4 + 4} dx = \frac{1}{2} \lim_{R \rightarrow \infty} \int_{-R}^R \frac{x^2}{x^4 + 4} dx$$

we will use Residue Th. for

$$\int_c \frac{z^2}{z^4 + 4} dz = \int_{c_1} \frac{z^2}{z^4 + 4} dz + \int_{c_2} \frac{z^2}{z^4 + 4} dz = 2\pi i \sum \text{Res at all poles inside } c$$

where  $c_1 : [-R, R]$  line segment and  $c_2 : |z| = R, \text{Im } z > 0$  half circle , for  $R > 2$

we can estimate

$$\left| \int_{c_2} \frac{z^2}{z^4 + 4} dz \right| \leq \max_{c_2} \left| \frac{z^2}{z^4 + 4} \right| \cdot \text{length } c_2 \leq \max_{|z|=R} \frac{|z|^2}{|z|^4 - 4} \cdot \pi R \leq \frac{\pi R^3}{R^4 - 4} \rightarrow 0$$

as  $R \rightarrow \infty$

Next, find all singularities of the integrand function i.e. solve  $z^4 = -4$

$$z^2 = \pm 2i \text{ and } z = \pm(1 + i) \text{ or } \pm(1 - i) \text{ OR } z = 4^{\frac{1}{4}} e^{i\frac{\pi}{4} + i\frac{\pi k}{2}} = \sqrt{2} e^{i\pi\left(\frac{1+2k}{4}\right)}$$

inside the curve only poles with  $\text{Im } z > 0$  so  $z_1 = 1 + i, z_2 = -1 + i$

$$\text{Res}[f, z_j] = \lim_{z \rightarrow z_j} (z - z_j) f(z) = \lim_{z \rightarrow z_j} (z - z_j) \frac{z^2}{z^4 + 4} = z_j^2 \cdot \frac{1}{4z_j^3} = \frac{1}{4z_j}$$

$$\text{Re } s [f, z_1] + \text{Re } s [f, z_2] = \frac{1}{4} \left[ \frac{1}{1+i} + \frac{1}{-1+i} \right] = -\frac{i}{4}$$

and the integral is  $2\pi i \cdot \frac{-i}{4} = \frac{\pi}{2}$  and  $\int_0^{\infty} \frac{x^2}{x^4 + 4} dx = \frac{\pi}{4}$ .

**For 9)**

$$\int_0^{2\pi} \frac{\cos 2\theta}{5 + 4 \cos \theta} d\theta = \operatorname{Re} \int_0^{2\pi} \frac{e^{i2\theta}}{5 + 4 \cos \theta} d\theta = \operatorname{Re} \oint_{|z|=1} \frac{z^2}{5 + 4 \left(\frac{z^2+1}{2z}\right)} \frac{dz}{iz} =$$

(by subst.  $z = e^{i\theta}$ ,  $\cos \theta = \frac{1}{2}(z + \frac{1}{z})$  and  $dz = ie^{i\theta} d\theta = iz d\theta$ ,  $e^{i2\theta} = z^2$ )

$$= \operatorname{Re} \frac{1}{i} \oint_{|z|=1} \frac{z^2}{5z + 2z^2 + 2} dz = \operatorname{Re} \frac{1}{i} \oint_{|z|=1} \frac{z^2}{2(z + \frac{1}{2})(z + 2)} dz =$$

$$= \operatorname{Re} \left[ \frac{1}{i} \cdot 2\pi i \cdot \operatorname{Res} \left[ f, -\frac{1}{2} \right] \right] = \frac{\pi}{6}$$

since only  $z = -\frac{1}{2}$  is inside the unit circle and

$$\operatorname{Res} \left[ f, -\frac{1}{2} \right] = \left[ \left( z + \frac{1}{2} \right) f(z) \right]_{z=-\frac{1}{2}} = \left[ \frac{z^2}{2(z+2)} \right]_{z=-\frac{1}{2}} = \frac{1}{12}.$$

**For 10a)**

the given branch of  $\log$  is cont. and therefore analytic in the  $z$ -plane except the branch cut :  $S = \left\{ \theta = \frac{\pi}{4} \text{ OR. } \operatorname{Im} z = \operatorname{Re} z \geq 0 \right\}$

so mapping is conformal on  $C - S$  since  $(\log z)' = \frac{1}{z}$  there.

**For b)**

for  $x = 0, y > 0$

$w = \ln y + i\frac{\pi}{2}$ .....horizontal line since  $u = \ln y$  is any real #,  $v = \frac{\pi}{2}$

for  $x = 0, y < 0$

$w = \ln(-y) + i\left(-\frac{\pi}{2} + 2\pi\right)$  .....horizontal line  $v = \frac{3\pi}{2}$

**For c)**

$|z| = 1$  means  $z = e^{i\theta}$  where  $\frac{\pi}{4} \leq \theta < \frac{9\pi}{4}$

$w = i\theta$ ... vert. line segment,  $u = 0, \frac{\pi}{4} \leq v < \frac{9\pi}{4}$